

Assessing the Effects of Submesoscale Parameterizations: Progress Report

Jody Klymak
School of Earth and Ocean Sciences
University of Victoria
P.O. Box 3055 STN CSC
Victoria, BC Canada, V8W 3P6
phone:(250)-472-5969 fax:(250)-721-6200 email:jklymak@uvic.ca
Award Number: N00014-08-1-1039
<http://web.uvic.ca/~jklymak/>



Figure 1: R/P FLIP deployed on the south of Monterey Bay during AESOP. The port boom, on the left, supported the Deep 8 Doppler system; the aft boom, in the foreground, supported the fastCTD. (Photos: T. Hughen)

LONG-TERM GOALS

Our long-term goal is to understand how energy is supplied to the ocean, and how it subsequently cascades to the turbulence and mixing important to the circulation, and the transport and distribution of tracers. This problem involves scales spanning subinertial motions to turbulence, and therefore requires integrative efforts with other sea-going investigators and numerical modelers. AESOP was an excellent opportunity to collect a data set in the coastal environment to test our understanding of the cascade.

OBJECTIVES

To test if high resolution models (i.e. SUNTANS, K. Winter's model) generate realistic internal waves due to tidal and wind forcing.

Report Documentation Page			Form Approved OMB No. 0704-0188		
Public reporting burden for the collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to a penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number.					
1. REPORT DATE 30 SEP 2008		2. REPORT TYPE		3. DATES COVERED 00-00-2008 to 00-00-2008	
4. TITLE AND SUBTITLE Assessing The Effects Of Submesoscale Parameterizations: Progress Report				5a. CONTRACT NUMBER	
				5b. GRANT NUMBER	
				5c. PROGRAM ELEMENT NUMBER	
6. AUTHOR(S)				5d. PROJECT NUMBER	
				5e. TASK NUMBER	
				5f. WORK UNIT NUMBER	
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) University of Victoria,School of Earth and Ocean Sciences,P.O. Box 3055 STN CSC,Victoria, BC Canada, V8W 3P6,				8. PERFORMING ORGANIZATION REPORT NUMBER	
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES)				10. SPONSOR/MONITOR'S ACRONYM(S)	
				11. SPONSOR/MONITOR'S REPORT NUMBER(S)	
12. DISTRIBUTION/AVAILABILITY STATEMENT Approved for public release; distribution unlimited					
13. SUPPLEMENTARY NOTES code 1 only					
14. ABSTRACT Our long-term goal is to understand how energy is supplied to the ocean, and how it subsequently cascades to the turbulence and mixing important to the circulation, and the transport and distribution of tracers. This problem involves scales spanning subinertial motions to turbulence, and therefore requires integrative efforts with other sea-going investigators and numerical modelers. AESOP was an excellent opportunity to collect a data set in the coastal environment to test our understanding of the cascade.					
15. SUBJECT TERMS					
16. SECURITY CLASSIFICATION OF:			17. LIMITATION OF ABSTRACT Same as Report (SAR)	18. NUMBER OF PAGES 5	19a. NAME OF RESPONSIBLE PERSON
a. REPORT unclassified	b. ABSTRACT unclassified	c. THIS PAGE unclassified			

To see if these models move energy from the large to the small scales properly.

To test how well Mellor-Yamada/KPP parameterizations do at estimating the location and magnitude of internal wave-driven mixing in these high resolution models.

To determine if the differences between observed and modeled mixing matter for quantities of operational and oceanographic relevance.

APPROACH

Progress is achieved through a cycle of instrument development, field observation, data analysis and comparison with theory and models. The primary instruments employed include Doppler sonar and rapidly profiling CTD's. Our instruments produce information that is quasi-continuous in space and time, typically spanning two decades in the wave number domain. This broad band space-time coverage enables the investigation of multi-scale interactions.

WORK COMPLETED

We deployed *R/P FLIP* off of Carmel CA, in 950 m of water for 18 days in Aug 2006. Data collected included velocity from the SIO Deep 8 Doppler Sonar and fastCTD, supplemented with data from a developmental wirewalker.

This data has been processed and supplied to other PIs in the program, particularly Sanford and Luther (for comparison with their EM sensors), Girton and Kunze (for comparison with their shipboard surveys) and Wang (model/data comparisons).

Higher-order processing has been started. This includes improved estimates of overturns for computing the turbulence dissipation rate, and assessing the feasibility of using the microconductivity sensor for estimates of scalar dissipations.

The purpose of this grant is to fund the analysis of this data and turn it into papers. We are only a month into the grant, so this phase of the work has just begun.

RESULTS

An example of the high resolution data set collected with the instruments deployed aboard *R/P FLIP* is shown in figure 2. The tide is clearly seen in the measurements both as alternating bands and as heaving in the isopycnals. Note how the lines of constant shear (shaded) follow the contoured isopycnals.

We have begun to use these measurements to characterize the wavefield in a number of ways. First, the observed energy flux at the site is relatively weak, and has an apparent spring-neap cycle (figure 3). The average number for the whole tidal cycle is 0.26 kW m^{-1} to the north and less than that to the east. Comparisons will be made to the energy fluxes in SUNTANS for similar forcing. Our measurements agree with Girton and Kunze's for the early part of our cruise when the tidal forcing was most similar.

We have also made progress in characterizing the turbulence and internal wave characteristics. Energy density ($KE = 0.5(u^2 + v^2)$ and $PE = 0.5(N^2\zeta^2)$) does not follow a clear spring-neap cycle, but rather appears to pick up during neap tide, particularly at mid depths (figure 4).




Figure 2: Velocity and density for 5 days during the cruise. Rapid sampling of the density helps deconvolve advection of velocity layers by displacements using the semi-Lagrange transform.

We have compared the dissipation rates observed using overturning scales and those estimated from the internal wave continuum energy levels using the Gregg-Henyey proxy (figure 5). The correspondence between the two is poor, particularly deeper than 400 m. Part of our efforts will be based on trying to improve these parameterizations, since their applicability appears limited except in the open ocean.

IMPACT/APPLICATIONS

Understanding the source of turbulence in the ocean is a primary goal. If we are successful, our understanding will lead to improved parameterizations for turbulence and mixing, which can be used in regional models.

RELATED PROJECTS

This data set is an important part of the AESOP effort, and we will be working closely with the other AESOP PIs to interpret and integrate it as described above.




Figure 3: Energy flux at the FLIP site for modes 1 and 2. Thick line is 1-day averages of the energy flux. No harmonic fits have been performed for these estimates.

This work is complementary to the work carried out in the Hawaiian Ocean Mixing Experiment, and we have already started comparisons with that data set. Similarly, ideas about the nature of the internal wave field that Pinkel developed for data collected as part of the Surface Heat Budget of the Arctic Ocean (SHEBA) experiment will be tested with this data.

Finally, it is complementary to Klymak's work at UVic, funded through the Canadian National Science and Engineering Research Council, to look at coastal internal wave processes. In particular, I have been investigating the impact of local breaking internal tides on mixing near topography.




Figure 4: Energy (KE and PE) observed at the R/P FLIP site.




Figure 5: Turbulence dissipation rate estimated at R/P FLIP using overturns (upper) and the Gregg-Henyey method (lower).